

A search for H_2 around pre-main-sequence stars

I. Sheret^{1*}, S. K. Ramsay Howat² and W. R. F. Dent²

¹*Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ*

²*UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ*

Accepted 1988 December 15. Received 1988 December 14; in original form 1988 October 11

ABSTRACT

We present the results of a search for pure rotational molecular hydrogen emission from two pre-main-sequence stars, AB Aur and CQ Tau. Observations were made using MICHELLE, the mid-IR echelle spectrometer at the UK Infrared Telescope. We found some evidence for emission in the $J = 4 \rightarrow 2$ line in AB Aur, but no $J = 3 \rightarrow 1$ line from either star. We derive upper limits on line flux which are significantly smaller than previous line flux estimates based on *ISO* observations. This suggests that the emission detected by *ISO* is extended on scales of at least 6 arcsec, and does not come from the disk as previously thought.

Key words: circumstellar matter.

1 INTRODUCTION

Circumstellar disks play a vital part in both star and planet formation, and if we are to understand these processes then it seems we must understand the composition and physics of disks first. The discovery that disks can survive well into the main sequence lifetime of a star (after planet formation ended in our own system) has opened up an exciting window on the planet formation process (Zuckerman 2001). This means that we can directly probe the environment of planet formation, even though planets themselves may be undetectable.

The dominant constituent of the disks around pre-main-sequence stars is H_2 . However, observing emission from H_2 which represents the bulk of the warm ($\sim 200\text{K}$) gas in the disks is not straightforward. Near-infrared (Bary et al. 2002), UV emission (Valenti et al. 2000) or absorption observations (Roberge et al. 2001) trace only the hottest gas or are dependent on the line of sight through the circumstellar disk. Mid-infrared observations of the pure rotational emission lines of H_2 trace the bulk of the warm gas, but are difficult from ground-based sites, due to telluric absorption and the high mid-infrared background emission. Thus, many previous studies of the gas component of such disks has relied on observations of tracer molecules, particularly CO (Mannings & Sargent 2000; Duvert et al. 2000). While these observations have been successful in tracing the velocity structure in disks, it seems that CO is a poor indicator of the total gas content. This is because CO can be destroyed by UV photons or frozen onto dust grains, leading to a severe underestimate of the total amount of gas. Most estimates of disk mass are based on sub-mm observations of

the dust continuum (Beckwith et al. 1990; Sylvester et al. 2001), but these rely on assumptions about the composition and size distribution of the dust grains, and also on the gas/dust mass ratio. More importantly, we know that the dust and gas do not evolve in the same way – disks around main-sequence stars are composed mainly of dust, while disks around pre-main-sequence stars are composed mainly of gas. It is important to be able to measure gas properties independently from the dust properties using direct observations of H_2 .

Recently, the Infrared Space Observatory (*ISO*) has been used to search for H_2 pure-rotational emission lines from pre-main-sequence stars, and also young main-sequence stars which are known to have dust disks. Thi et al. (2001b) report detections of the H_2 $J = 3 \rightarrow 1$ transition ($\lambda = 17.035 \mu\text{m}$) and H_2 $J = 2 \rightarrow 0$ transition ($\lambda = 28.221 \mu\text{m}$). H_2 masses derived from these observations imply large masses of H_2 in the disks around main-sequence targets (Thi et al. 2001a), reversing the previous belief that dust is the main component. However, as the *ISO* beam is large (14×27 arcsec) compared with the disks and the observations have moderate spectral resolution ($R \sim 2000$), these results cannot distinguish circumstellar disk emission from that of a foreground cloud or extended source.

Follow up observations of both the pure rotational emission lines and H_2 in absorption have provided strong evidence that the warm H_2 is extended in sources selected from the *ISO* studies. Using the the echelle spectrometer, TEXES, Richter et al. (2002) obtained observations of six sources, three of which have reported *ISO* H_2 detections (Thi et al. 2001b): HD163296, AB Aur and GG Tau. They obtained upper limits on the line fluxes from five sources, with a possible 2 sigma detection of the H_2 $J = 3 \rightarrow 1$ S(0) line in AB Aur and conclude that the majority of the H_2

* E-mail: is@roe.ac.uk

emission detection by *ISO* is extended on scales of 5 arcsec or more. The far-UV spectrum of Beta Pictoris presented by Lecavelier des Etangs et al. (2001) shows a complete absence of H_2 absorption lines, contrary to expectations if large quantities of H_2 are present within the edge-on dust disk.

We present the results of our attempts to confirm the *ISO* results using MICHELLE, the mid-infrared echelle spectrometer on the ground-based United Kingdom Infrared Telescope (UKIRT). We targeted two pre-main-sequence stars, CQ Tau (distance 100 pc) and AB Aur (distance 142 pc), both of which have good *ISO* detections and one of which (AB Aur) was also observed by Richter et al. (2002). In both cases, the dust disk is expected to be unresolved spatially by MICHELLE.

2 OBSERVATIONS AND DATA REDUCTION

Mid-IR echelle spectroscopy of AB Aur and CQ Tau was carried out on 2001 December 29 and 30 using MICHELLE (Glasse et al. 1997) on UKIRT. AB Aur and CQ Tau were observed in the H_2 $J = 3 \rightarrow 1$ transition ($\lambda = 17.035 \mu\text{m}$), providing direct comparison with the *ISO* observations. AB Aur was also observed in the H_2 $J = 4 \rightarrow 2$ transition ($\lambda = 12.279 \mu\text{m}$). The H_2 $J = 2 \rightarrow 0$ transition ($\lambda = 28.221 \mu\text{m}$) was used in the *ISO* observations, but is not readily accessible from the ground due to the structure of the atmospheric transmission at $28 \mu\text{m}$. Details of the observations are shown in Table 1.

MICHELLE is a mid-IR imager and spectrometer which uses a 320×240 -pixel Si:As array and operates between 8 and 25 microns. For these observations the echelle grating was used, to provide maximum sensitivity to the H_2 lines which are expected to be unresolved. AB Aur was observed with a 2-pixel slit width; CQ Tau with a 4-pixel slit width. The resolving power was $R = 15\,200$ at $12 \mu\text{m}$ and $R = 18\,600$ ($R = 9\,300$) at $17 \mu\text{m}$ for a two (four) pixel wide slit; as a result the H_2 lines are well separated from the bright sky lines. A single order from the grating is selected using blocking filters. The pixel field of view varies with wavelength due to anamorphic demagnification. The values are given in Table 1.

The point-like sources were all observed by nodding along the 70 arcsec long slit, which was orientated north-south on the sky. The separation of the two beams was 15 arcsec and the dwell-time in each beam was 40 seconds.

Data reduction was done using software written in the IDL programming language. Pairs of frames from different nod positions were subtracted, giving a set of difference frames each containing a positive and a negative version of the source. In each frame, the positive and negative versions were combined to give a single 2D spectrum. Before the set of 2D spectra were combined into a final image, the dataset was despiked using an iterative sigma cut. To do this, all of the 2D spectra were compared, and datum were flagged as bad if they were further than a specified number of standard deviations from the mean of their pixel. This sigma cut was done three times with threshold settings of 6, 4 and 3 sigma.

Once a final 2D spectrum had been produced, 1D spectra were extracted in two ways: a wide extraction which summed all the emission in a 6.5 arcsec region centered on the source, and an optimal extraction of the continuum. The




Figure 1. Spectra of the H_2 $J = 3 \rightarrow 1$ transition for AB Aur and CQ Tau. For comparison, gaussians are shown with the same integrated flux as the *ISO* measurements reported in Thi et al. (2001b), assuming an intrinsic line width of 20 km s^{-1} . The CQ Tau spectrum has been masked out at regions of high noise (shown by the use of dotted lines); the lower panel shows the standard star observation, scaled from 0.0 to 1.0, to demonstrate that the masked regions correspond to atmospheric absorption lines. For AB Aur, the standard star observation was saturated, which causes the variability in the continuum level.

optimal extraction method takes a weighted mean of the pixels around the source, giving more weight where the continuum is brighter. This method gives the lowest possible noise on the continuum level, but if the H_2 emission region is more extended than the continuum emission then it will underestimate the H_2 line flux. For both extraction methods a sky subtraction was done first, estimating the residual sky signal in each column using pixels between 3.5 and 7 arcsec from the source.

Standard stars were observed and reduced in the the same way as the science targets. The final spectrum for each object was divided through by a standard spectrum to remove wavelength dependant variations in atmospheric and instrumental transmission. The standard stars were ob-

Object	Date	Wavelength μm	Slit width pixels	Pixel scale on-sky arcsec	Integration time seconds
AB Aur	29 Dec 2001	12.279	2	0.59×0.24	1900
	29 Dec 2001	17.035	2	0.50×0.28	3400
CQ Tau	30 Dec 2001	17.035	4	0.50×0.28	5900

Table 1. Summary of observations. The pixel scale on the sky is subject to anamorphic magnification due to the quasi-Littrow mode of operation of the MICHELLE spectrometer. The long axis is along the slit.

Object	Wavelength μm	Line flux $\times 10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2}$		
		Wide extraction	Optimal extraction	<i>ISO</i> Result
AB Aur	12.297	5.4 ± 3.8	4.8 ± 2.1	
	17.035	< 14.6	< 8.9	30 ± 9
CQ Tau	17.035	< 13.0	< 6.7	40 ± 12

Table 2. Summary of results. All upper limits are 3 sigma. *ISO* results are taken from Thi et al. (2001b).

served using the same rows on the detector as the sources, and division by the standard effectively flat-fields the detector. For the 17 μm observations of AB Aur the standard star observation was saturated, so we were unable to make this correction. This causes variation in the continuum level, but would not prevent us from detecting a narrow emission line.

Wavelength calibration was obtained by measuring the positions of sky lines, and comparing the positions to a model sky spectrum. We estimate a 1 sigma error on the wavelength calibration is about one pixel ($\sim 3 \times 10^{-4} \mu\text{m}$), and is largely due to the difficulty in centroiding the broad sky lines. The spectra were shifted to the object's rest frame, accounting for the radial velocity of the earth and object's heliocentric radial velocity ($+8.9 \text{ km s}^{-1}$ for AB Aur and $+22.6 \text{ km s}^{-1}$ for CQ Tau (Barbier-Brossat & Figon 1999)). Flux calibration was based on the observed continuum level and published continuum flux measurements from Thi et al. (2001b) for both the 17 μm spectra and Richter et al. (2002) for the AB Aur 12 μm spectrum.

3 RESULTS

Figures 1 and 2 show the final spectra produced using an optimal extraction. At the location of sky lines the noise increases dramatically, so these regions have been masked out (i.e. they were not used when estimating the continuum level and the overall noise). On the 17 μm figures a gaussian with the total line flux measured by Thi et al. (2001b) is shown, to allow comparison between the *ISO* results and our spectra. As the *ISO* observations do not resolve the line, we have assumed an intrinsic line width of 20 km s^{-1} (see Section 4 for discussion of this value), and an instrumental width of 16 km s^{-1} for AB Aur and 32 km s^{-1} for CQ Tau, giving a total width of 26 km s^{-1} and 38 km s^{-1} respectively. The MICHELLE sensitivities should have produced a clear detection of the line, if all of the flux in the $14 \times 27 \text{ arcsec}$ *ISO* beam were contained within our extracted beams, as expected for emission from an unresolved disk.

To estimate the line flux, all of the flux within a 40 km s^{-1} wide bin centered on the systematic velocity was




Figure 2. Spectra of the H₂ $J = 4 \rightarrow 2$ transition for AB Aur. The vertical dotted line shows the expected location of the transition. Noisy regions have been masked out and the standard star is shown in the lower panel, as in Figure 1.

summed, after subtracting the continuum level. Fluxes and upper limits for all extraction methods are shown in Table 2. There is no evidence of H₂ emission in either of the 17 μm spectra. The 12 μm optimally extracted spectrum gives flux of $4.8 \pm 2.1 \text{ ergs s}^{-1} \text{ cm}^{-2}$, but as this is only a 2.3 sigma result should not be regarded as a definite detection. However, this result is consistent with the tentative detection by Richter et al. (2002), who measure $(2.0 \pm 1.0) \times 10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2}$ with a width of 10.5 km s^{-1} .

4 DISCUSSION

Despite the enhanced sensitivity to narrow emission lines from a point-like source, these new spectra of AB Aur and CQ Tau show no convincing detections of either the 0-0 S(1)

or 0-0 S(2) lines. In order for the 0-0 S(1) H₂ emission to be detectable by *ISO* but not by MICHELLE, the emission would have to be either spectrally broad or spatially extended. We consider these two explanations in turn.

The spectral resolving power of the *ISO* observations is $R (= \frac{\lambda}{\delta\lambda}) = 2400$, and the H₂ lines Thi et al. (2001b) observed are unresolved. In calculating our upper limits, we assume that the lines had an intrinsic width of 20 km s⁻¹. The dominant line broadening mechanisms for gas in a disk are thermal broadening, and bulk motion due to the disk's rotation. Assuming the gas has a temperature of 200 K (as estimated by Thi et al. (2001b)), the thermal broadening would only be 1.5 km s⁻¹. The amount of rotational broadening depends on the distance from the star, and the inclination of the disk. If we assume a distance of 2AU (a reasonable choice, given the gas temperature of 200 K), this gives a width of 15 km s⁻¹, assuming an inclination of $\sim 45^\circ$. Therefore, 20 km s⁻¹ is a conservative estimate of the line width. In fact, the line FWHM would have to be over 100 km s⁻¹ to make the *ISO* and MICHELLE measurements consistent.

The possibility of extended emission, due to excitation of a remnant cloud of molecular gas by UV photons or by shocks, is plausible. Assuming that the *ISO* measured flux is uniformly distributed over the 14×27 arcsec aperture, the fluxes are consistent with UV-excited emission strengths predicted by Burton et al. (1992) for moderate UV field strengths ($10^3 - 10^4$ times that of the interstellar medium) and moderate densities ($n_{\text{H}_2} \sim 10^3 - 10^4 \text{ cm}^{-3}$). Over the MICHELLE beam used to obtain the spectra shown in Figure 1, the line flux would be an order of magnitude fainter than the sensitivity limit obtained. Moreover, the sky subtraction region (3.5-7 arcsec radius) means that our observations are insensitive to emission at radius > 3.5 arcsec. The hypothesis of extended emission is strengthened for AB Aur since a map of ¹²CO 3-2 shows an extended molecular cloud around the central source, with a size of at least 30 arcsec (Thi et al. 2001b). For CQ Tau, no ¹²CO 3-2 line was detected, but this could be reconciled with extended H₂ emission if the CO has been destroyed by photodissociation (Dent et al. 1995).

In interpreting our results, an important question concerns the optical depth of the dust disk. If the dust is optically thick at the wavelengths we observed and the gas and dust are at the same temperature, then the gas would be equally likely to absorb or emit a photon, so no emission line would be produced. Line emission will only occur if there are gaps in the dust disk, or if there is a hot atmosphere above the disk. Though these situations are quite plausible, only a small fraction of the gas in the disk would be observed in this case, and any calculation of the gas mass would underestimate the true mass. It is therefore essential that we know the optical depth of the disk in order to correctly interpret our results.

The optical depth of the disks can be estimated from radiative transfer models based on the SED of the disks. Natta et al. (2001) made models of the disk SEDs for both AB Aur and CQ Tau, and we can use their fitted parameters to determine whether the disks are optically thick at 12 and 17 μm . Based on their estimates of disk mass and disk radius, and using the usual assumption that the dust opacity at 1.3 mm is $\kappa_{1.3\text{mm}} = 0.01 \text{ cm}^2 \text{ g}^{-1}$, the average

optical depth of the disks at 1.3 mm is 0.03 for AB Aur and 0.24 for CQ Tau. Then, using a power law dust opacity index ($\kappa \propto \lambda^{-\beta}$, where β was estimated to be 2.0 for AB Aur and 1.0 for CQ Tau in Natta et al. 2001), the optical depth can be estimated at shorter wavelengths. This indicates that the disks become optically thick at wavelengths shorter than 200 μm for AB Aur and 300 μm for CQ Tau. Both disks must therefore have a high optical depth at 12 and 17 μm .

This result has two important implications. Firstly, this means that we cannot use our upper limits on line flux to calculate upper limits on the mass of H₂ in the disk, as most of the gas in the disk is hidden by the optically thick dust. The H₂ line at 12 μm (if confirmed) must arise from a hot atmosphere above the disk, or from gaps in the dust disk. Secondly, if the lines detected by *ISO* do arise in the disk, then only a small fraction of the gas was observed and the total gas mass must be much higher than was estimated.

In conclusion, we have searched for pure rotational H₂ emission lines from the circumstellar disks of AB Aur and CQ Tau. We tentatively detect emission from AB Aur, but find no evidence for emission from CQ Tau. The likely reason for this non-detection is that the disks are optically thick in the mid-IR. Our upper limits on line flux are significantly smaller than previous line flux estimates based on *ISO* observations, which suggests that the emission detected by *ISO* is extended on scales of at least 6 arcsec (or $> 60\text{AU}$).

Mapping the emission over the area covered of the *ISO* beam is one approach to confirming these conclusions. Detecting the faint extended emission will require a considerable gain in sensitivity over these observations, such as is anticipated using MICHELLE on the Gemini North 8-m telescope.

The United Kingdom Infrared Telescope is operated by the Joint Astronomy Centre on behalf of the UK Particle Physics and Astronomy Research Council. We wish to thank the staff at UKIRT for enabling these observations. We also thank Mark Wyatt and Alistair Glasse for helpful discussions of this work.

REFERENCES

- Barbier-Brossat M., Figon P., 1999, VizieR Online Data Catalog, 3213, 0
- Bary J. S., Weintraub D. A., Kastner J. H., 2002, ApJ, 576, L73
- Beckwith S. V. W., Sargent A. I., Chini R. S., Guesten R., 1990, AJ, 99, 924
- Burton M. G., Hollenbach D. J., Tielens A. G. G., 1992, ApJ, 399, 563
- Dent W. R. F., Greaves J. S., Mannings V., Coulson I. M., Walther D. M., 1995, MNRAS, 277, L25
- Duvert G., Guilloteau S., Ménard F., Simon M., Dutrey A., 2000, A&A, 355, 165
- Glasse A. C., Atad-Ettinger E. I., Harris J. W., 1997, in Proc. SPIE Vol. 2871, p. 1197-1203, Optical Telescopes of Today and Tomorrow, Arne L. Ardeberg; Ed., Vol. 2871, pp. 1197-1203

- Lecavelier des Etangs A., Vidal-Madjar A., Roberge A., Feldman P. D., Deleuil M., André M., Blair W. P., Bouret J.-C., Désert J.-M., Ferlet R., Friedman S., Hébrard G., Lemoine M., Moos H. W., 2001, *Nature*, 412, 706
- Mannings V., Sargent A. I., 2000, *ApJ*, 529, 391
- Natta A., Prusti T., Neri R., Wooden D., Grinin V. P., Mannings V., 2001, *A&A*, 371, 186
- Richter M. J., Jaffe D. T., Blake G. A., Lacy J. H., 2002, *ApJ*, 572, L161
- Roberge A., Lecavelier des Etangs A., Grady C. A., Vidal-Madjar A., Bouret J.-C., Feldman P. D., Deleuil M., André M., Boggess A., Bruhweiler F. C., Ferlet R., Woodgate B., 2001, *ApJ*, 551, L97
- Sylvester R. J., Dunkin S. K., Barlow M. J., 2001, *MNRAS*, 327, 133
- Thi W. F., Blake G. A., van Dishoeck E. F., van Zadelhoff G. J., Horn J. M. M., Becklin E. E., Mannings V., Sargent A. I., van den Ancker M. E., Natta A., 2001a, *Nature*, 409, 60
- Thi W. F., van Dishoeck E. F., Blake G. A., van Zadelhoff G. J., Horn J., Becklin E. E., Mannings V., Sargent A. I., van den Ancker M. E., Natta A., Kessler J., 2001b, *ApJ*, 561, 1074
- Valenti J. A., Johns-Krull C. M., Linsky J. L., 2000, *ApJS*, 129, 399
- Zuckerman B., 2001, *ARA&A*, 39, 549

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.